

COLORADO GEOLOGICAL SURVEY
Open-file Report OF-18-10
Debris Flow Susceptibility Map of Jefferson County, Colorado
Kevin M. McCoy, Matthew L. Morgan, and Karen A. Berry

ABOUT THIS MAP

This map is intended for use by planners and regulators to support review of site-specific geologic hazard reports submitted for development purposes as required by law, and by professional geologists planning detailed site-specific geologic hazard studies. The red polygons represent areas that may be susceptible to debris flows, especially during extreme precipitation events. New and existing structures, roadways, bridges, utilities, and other infrastructure located within these mapped debris-flow susceptible areas may be at risk of structural damage and/or sediment inundation. For new/proposed development within these areas, site-specific geologic hazard reports should be required prior to approval of land subdivision or the issuance of building permits. These reports should discuss the degree, limits, and potential impacts of the hazard to the proposed development or land use changes; proper debris-flow mitigation techniques; and feasibility of any recommended mitigation techniques.

DESCRIPTION OF THE HAZARD

For this map and report, the term “debris flow” comprises a range of related natural phenomena variously described in Colorado State statutes (Rogers et al., 1974) and scientific and engineering literature (e.g., Costa, 1988; Pierson, 2005; Moase et al., 2017) as mudflows, debris flows, and/or sediment-laden floods (hyperconcentrated flows). Distinguishing between these phenomena can be important for site-specific hazard analysis and mitigation design, but is beyond the scope of this project.

Debris flows are common hazards in mountain and piedmont valleys and along the steep flanks of buttes, mesas, and hogbacks in Colorado (e.g., Mears, 1977; Costa and Jarrett, 1981; Coe and Godt, 2003; Coe et al., 2003; Godt and Savage, 2003; Morgan et al., 2004; Godt and Coe, 2007; Coe et al., 2014). They typically form when heavy rainfall or rapid snowmelt events trigger landslides on steep slopes and/or flash floods in steep drainages. Intense wildfires can increase the likelihood of debris flow initiation by removing soil-stabilizing vegetation and reducing the soil’s ability to absorb rainfall and rapid snowmelt; as a result, relatively common (e.g. 2-year recurrence) rain storms can trigger debris flows in burned areas where debris-flow initiation would typically require a relatively rare extreme rainfall or rapid snowmelt event. In general, the risk of debris-flow initiation from relatively common rain storms tends to be greatest in the first few years after a major fire; however, elevated debris flow hazard in burned basins can persist for a period of time ranging from a few years to several decades or more, until the vegetation recovers and soil infiltration returns to pre-fire conditions or sufficient erodible material is removed from the basin.

Once initiated, debris flows can grow rapidly, transforming into muddy, sediment-charged slurries that can become as dense as wet concrete as they erode hillslope soils and channel sediment and entrain loose debris. Fully formed debris flows can transport rocks, trees, and other debris for long distances until the channel eventually widens and/or the channel slope flattens and the entrained material begins to deposit, usually on an alluvial and/or debris fan. Alluvial/debris fan flooding can be much more spatially random than typical floodplain flooding along streams; it is typically characterized by lateral movement of flow over a wide area from the mouth of a drainage basin onto a lower lying area. Established channels can suddenly fill with sediment and debris, be abandoned, and new channels formed during a single event. For that reason, structures and infrastructure developed on alluvial or debris fans may be susceptible to debris-flow impacts, even if they are located relatively far from the established stream channel.

Historically, human development along mountain fronts and within mountain and piedmont valleys commonly occurred on alluvial fans. Roads and other linear infrastructure in these areas may cross alluvial fans or debris-flow generating drainages. Debris flows that reach and inundate structures or other manmade features in these areas may cause serious issues including: injury or death, potentially costly outlay of funds for repairs and (or) sediment removal, temporary road closures, and temporary or permanent loss of access to buildings, roads, and other infrastructure. Debris flows can block

improperly designed culverts where roads cross debris fans or steep drainages, leading to unanticipated flooding, deposition of debris on roadways, damage to the road surfaces, or washout of the road. Where debris-flow generating tributaries meet larger streams, deposition of entrained material can lead to upstream flooding due in part to temporary damming and downstream aggradation. The latter commonly results in degraded water quality owing to increased sediment loading.

MAPPING METHODOLOGY

This map was prepared by combining GIS-based debris-flow source and runout models with visual interpretation of high-resolution lidar-based digital terrain data, digital geologic maps, and digital soil survey data. The lidar data used has a nominal horizontal accuracy of 1-m and is publicly available for the entire county on the USGS National Map viewer (<https://viewer.nationalmap.gov/basic/>). Burned area outlines for fires that have occurred since the year 2000 were delineated using the US Historic Fire Perimeters dataset downloaded from the Geospatial Multi-Agency Coordination (GEOMAC) website (<https://www.geomac.gov/>). The areal extent of the 1996 Buffalo Creek fire was roughly estimated based on vegetation changes still visible in recent aerial imagery. Figures 1 and 2 show the coverage of digital geologic maps and digital soil survey maps used for this study, respectively.

Debris-flow source areas, typically defined by steep gullies and colluvial hollows where potentially debris-flow generating landslides, erosion, and transport may occur, were automatically identified in ArcGIS by overlaying slope, plan curvature, and flow accumulation rasters derived from a lidar-based digital elevation model (DEM). The automatically identified source areas were evaluated and manually revised by the analyst to exclude source areas considered to be erroneously generated and to include potential source areas missed by the automated model. Debris-flow runout zones were modeled using the MATLAB-based runout modeling program Flow-R (<http://www.flow-r.org/>). The resulting polygons were evaluated and manually revised by the analyst to produce the final susceptibility polygons shown on the map.




Figure 1. Index of geologic map data sources available for the study area.



Figure 2. *Index of digital soil survey reports available for the study area.*

HISTORICAL DEBRIS FLOW EVENTS IN THE COUNTY

An extreme rainfall event between September 9 – 13, 2013 caused deadly floods and landslides, including at least 1,138 debris flows, in Boulder, Larimer, and Jefferson counties (Coe et al., 2014). Three of the eight reported fatalities from the event were attributed to debris flows in Boulder County (Godt et al. 2014). Post-event landslide and debris flow mapping was performed by Morgan et al. (2013a, b). Coe et al. (2014), Godt et al. (2014), and Anderson et al. (2015), discuss conditions that led to debris flow initiation during the event.

Six named fires have burned various mountainous portions of Jefferson County since 1996:

1996 Buffalo Creek Fire
2000 High Meadow Fire
2002 Hayman Fire
2011 Indian Gulch Fire
2012 Lower North Fork Fire
2013 Lime Gulch Fire

Burn scars from the High Meadow, Buffalo Creek, and Hayman fires are still clearly visible in recent aerial imagery. Apparent debris flow/debris flood scars and deposits in these burned areas are visible in post-fire aerial imagery and in the LiDAR-based DEM. Post-fire hazard assessments and results of special studies for most of these fires are discussed by Moody and Martin (2001), Graham (2003), Stevens (2013), Elliott and Parker, (2001), Elliott et al. (2005), Ruddy (2011), and Jefferson Conservation District.

MAP LIMITATIONS

This map depicts generalized areas that may be susceptible to debris flows based on available GIS data and limited field observations in easily accessible areas. The map was generated at a scale of approximately 1:24,000 (1 inch = approx. 0.4 mi.) and is not valid if enlarged to scales greater than 1:24,000.

The degree of susceptibility to a particular geologic hazard, in any given area, is related to ever-changing natural and human-induced conditions, and any alteration in the natural landscape may increase or decrease susceptibility to a particular hazard. The polygons shown on this map are not intended to assign risk, or indicate the degree, severity, recurrence interval, or exact boundary of individual debris flows or debris-flow susceptible areas. Because of limitations associated with the map scale and the scope of this project, some areas that may be susceptible to potentially damaging but localized debris flows near the mouths of small gullies or at the base of steep slopes may not be included. Additionally, some higher-order streams in larger mountain valleys that have been mapped as debris flow susceptibility areas may be more likely to produce water floods than debris flows depending on local rainfall-runoff conditions and availability of erodible sediment and debris in the basin. Inclusion of existing structures and infrastructure within a mapped susceptibility area does not necessarily indicate that debris-flow impacts will occur, it only indicates that these features may be more exposed to debris-flow events than similar features located in other areas. This map should not be used in place of a detailed site-specific geologic hazard study.

ACKNOWLEDGEMENTS

The authors would like to thank Dr. Ralph Shroba for his review, comments, and suggestions, which improved the map and report. Additional lidar data was collected for this study with partial funding support from the Colorado Water Conservation Board.

REFERENCES CITED

- Anderson, S.W., Anderson, S.P., and Anderson, R.S., 2015, Exhumation by debris flows in the 2013 Colorado Front Range storm: *Geology*, v. 43, p. 391–394, doi: 10.1130/G36507.1.
- Bryant, B., 1976, Reconnaissance geologic map of the Bailey quadrangle, Jefferson and Park Counties, Colorado: United States Geological Survey Miscellaneous Geologic Investigations Map MF-816.
- Bryant, B., 1974a, Reconnaissance geologic map of the Conifer quadrangle, Jefferson County, Colorado: United States Geological Survey Miscellaneous Field Studies Map MF-597.
- Bryant, B., 1974b, Reconnaissance geologic map of the Pine quadrangle, Jefferson County, Colorado: United States Geological Survey Miscellaneous Field Studies Map MF-598.
- Bryant, B., Miller, R.D., and Scott, G.R., 1973, Geologic map of the Indian Hills quadrangle, Jefferson County, Colorado: United States Geological Survey Geologic Quadrangle Map GQ-1073.

- Coe, J.A., and Godt, J.W., 1999, Historical Debris Flows Along the Interstate-70 Corridor in Clear Creek County, Central Colorado: United States Geological Survey.
- Coe, J.A., Godt, J.W., Parise, M., and Moscariello, A., 2003, Estimating debris-flow probability using fan stratigraphy, historic records, and drainage-basin morphology, Interstate 70 highway corridor, central Colorado, U.S.A.
- Coe, J.A., Kean, J.W., Godt, J.W., Baum, R.L., Jones, E.S., Gochis, D.J., and Anderson, G.S., 2014, New insights into debris-flow hazards from an extraordinary event in the Colorado Front Range: *GSA Today*, v. 24, p. 4–10, doi: 10.1130/GSATG214A.1.
- Costa, J.E., 1988, Rheologic, geomorphic, and sedimentologic differentiation of water floods, hyperconcentrated flows, and debris flows, *in* Baker, V.R., Kochel, C.R., and Patton, P.C. eds., *Flood geomorphology*, p. 113–122.
- Costa, J.E., and Jarrett, R.D., 1981, Debris Flows in Small Mountain Stream Channels of Colorado and Their Hydrologic Implications: *Bulletin of the Association of Engineering Geologists*, v. XVIII, p. 309–322.
- Elliott, J.G., and Parker, R.S., 2001, Developing a post-fire flood chronology and recurrence probability from alluvial stratigraphy in the Buffalo Creek watershed, Colorado, USA: *Hydrological Processes*, v. 15, p. 3039–3051, doi: 10.1002/hyp.390.
- Elliott, J.G., Smith, M.E., Friedel, M.J., Stevens, M.R., Bossong, C.R., Litke, D.W., Parker, R.S., Costello, C., Wagner, J., Char, S.J., Bauer, M.A., and Wilds, S.R., 2005, Analysis and mapping of post-fire hydrologic hazards for the 2002 Hayman, Coal Seam, and Missionary Ridge wildfires, Colorado: USGS Scientific Investigations Report 2004–5300, <https://pubs.er.usgs.gov/publication/sir20045300>.
- Gable, D.J., 1972, Geologic map of the Tungsten quadrangle, Boulder, Gilpin, and Jefferson Counties, Colorado: United States Geological Survey Geologic Quadrangle Map GQ-978.
- Godt, J.W., and Savage, J.E., 2003, Geomorphometry of Small Debris-Flow Drainage Basins Along Interstate 70, Eagle County, Colorado, *in* Boyer, D.D., Santi, P.M., and Rogers, W.P. eds., *Engineering Geology in Colorado, Contributions, Trends, and Case Histories*, Association of Engineering Geologists Special Publication No. 15, Colorado Geological Survey Special Publication 55.
- Godt, J.W., and Coe, J.A., 2007, Alpine debris flows triggered by a 28 July 1999 thunderstorm in the central Front Range, Colorado: *Geomorphology*, v. 84, p. 80–97, doi: 10.1016/j.geomorph.2006.07.009.
- Godt, J.W., Coe, J.A., Kean, J.W., Baum, R.L., Jones, E.S., Harp, E.L., Staley, D.M., and Barnhart, W.D., 2014, Landslides in the Northern Colorado Front Range Caused by Rainfall, September 11–13, 2013: Fact Sheet 2013-3114.
- Graham, R.T., 2003, Hayman Fire Case Study: US Department of Agriculture, Forest Service, Rocky Mountain Research Station General Technical Report RMRS-GTR-114, https://www.fs.fed.us/rm/pubs/rmrs_gtr114.pdf.
- Jefferson Conservation District Land Rehabilitation FAQ Lower North Fork Fire: <http://www.jeffersonconservationdistrict.org/wp-content/uploads/2012/04/LNF-LandRehabFAQ.pdf>.
- Lindvall, R.M., 1978, Geologic Map of the Fort Logan quadrangle, Jefferson, Denver, and Arapahoe Counties, Colorado: USGS Geologic Quadrangle Map GQ-1427.
- Lindvall, R.M., 1979, Geologic map of the Arvada quadrangle, Adams, Denver, and Jefferson Counties, Colorado: USGS Geologic Quadrangle Map GQ-1453.
- Machette, M.N., 1977, Geologic map of the Lafayette quadrangle, Adams, Boulder, and Jefferson Counties, Colorado: United States Geological Survey Geologic Quadrangle Map GQ-1392.
- Mears, A.I., 1977, Debris-Flow Hazard Analysis and Mitigation; An Example from Glenwood Springs, Colorado:

Colorado Geological Survey Information Series IS-08.

- Moase, E., Strouth, A., Jakob, M., and Stead, D., 2017, Guidance for Steep Creek Hazard Mitigation Design in Canada, *in* Ottawa, Ontario, Canada.
- Moody, J.A., and Martin, D.A., 2001, Hydrologic and Sedimentologic Response of Two Burned Watersheds in Colorado: United States Geological Survey Water Resources Investigative Report 01-4122, <https://pubs.usgs.gov/wri/2001/4122/report.pdf>.
- Morgan, M.L., Temple, J., Grizzell, M.T., and Barkmann, P.E., 2004, Geologic Map of the Dawson Butte Quadrangle, Douglas County, Colorado: Colorado Geological Survey Open-File Report OF-04-7.
- Morgan, M.L., Fitzgerald, F.S., and Morgan, K.S., 2013a, Preliminary Survey of Debris Flow, Landslide, and Rockfall Deposits as a result of the September 11-14, 2013 Flooding Events, Boulder County, Colorado: <http://www.arcgis.com/home/item.html?id=39e6c721635f40c8add90112c9d1a646>.
- Morgan, M.L., Fitzgerald, F.S., and Morgan, K.S., 2013b, Preliminary Survey of Debris Flow, Landslide, and Rockfall Deposits as a result of the September 11-14, 2013 Flooding Events, Larimer County, Colorado: <http://www.arcgis.com/home/item.html?id=47fef299bc4a4a4c9e19cbb8afe66f4>.
- Peterson, W.L., 1964, Geology of the Platte Canyon quadrangle, Colorado: United States Geological Survey Bulletin B-1181-C.
- Pierson, T.C., 2005, Distinguishing between debris flows and floods from field evidence in small watersheds: United States Geological Survey Fact Sheet 2004-3142, <http://pubs.usgs.gov/fs/2004/3142/>.
- Rogers, W.P., Ladwig, L.R., Hornbaker, A.L., Schwochow, S.D., Hart, S.S., Shelton, D.C., Scroggs, D.L., and Soule, J.M., 1974, Guidelines and Criteria for Identification and Land-Use Controls of Geologic Hazard and Mineral Resource Areas: Colorado Geological Survey Special Report SP-06.
- Ruddy, B.C., 2011, Probability and Volume of Potential Postwildfire Debris Flows in the 2011 Indian Gulch Burn Area, near Golden, Colorado: United States Geological Survey Open-File Report 2011-1248, <https://pubs.usgs.gov/of/2011/1248/OF11-1248.pdf>.
- Scott, G.R., 1972, Geologic map of the Morrison quadrangle, Jefferson County, Colorado: United States Geological Survey Miscellaneous Geologic Investigations Map I-790-A.
- Scott, G.R., 1962, Geology of the Littleton quadrangle, Jefferson, Douglas, and Arapahoe Counties, Colorado: United States Geological Survey Bulletin B-1121-L.
- Scott, G.R., 1963, Quaternary geology and geomorphic history of the Kassler quadrangle, Colorado: U.S. Geological Survey Professional Paper PP-421-A.
- Sheridan, D.M., and Marsh, S.P., 1976, Geologic map of the Squaw Pass quadrangle, Clear Creek, Jefferson, and Gilpin Counties, Colorado: United States Geological Survey Geologic Quadrangle Map GQ-1337.
- Sheridan, D.M., Maxwell, C.H., Albee, A.L., and Van Horn, R., 1958, Preliminary Map of Bedrock Geology of the Ralston Buttes Quadrangle, Jefferson County, Colorado: United States Geological Survey Mineral Investigations Field Studies Map MF-179.
- Sheridan, D.M., Reed, J.C., and Bryant, B., 1972, Geologic map of the Evergreen quadrangle, Jefferson County, Colorado: United States Geological Survey Miscellaneous Geologic Investigations Map I-786-A.
- Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. Soil Survey Geographic (SSURGO) database for Golden Area, Colorado, Parts of Denver, Douglas, Jefferson, and Park Counties. Available online.

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. Soil Survey Geographic (SSURGO) database for Pike and San Isabel NF, Colorado, Northern Part, Parts of Chaffee, Clear Creek, Fremont, Jefferson, Lake, Park and Saguache Counties. Available online.

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. Soil Survey Geographic (SSURGO) database for Pike National Forest, Eastern Part, Colorado, Parts of Douglas, El Paso, Jefferson, and Teller Counties. Available online.

Spencer, F.D., 1961, Bedrock geology of the Louisville quadrangle, Colorado: United States Geological Survey Geologic Quadrangle Map GQ-151.

Stevens, M.R., 2013, Analysis of Postfire Hydrology, Water Quality, and Sediment Transport for Selected Streams in Areas of the 2002 Hayman and Hinman Fires, Colorado: United States Geological Survey Scientific Investigations Report 2012–5267, <https://pubs.usgs.gov/sir/2012/5267/>.

Taylor, R.B., 1976, Geologic map of the Black Hawk quadrangle, Gilpin, Jefferson, and Clear Creek Counties, Colorado: United States Geological Survey Geologic Quadrangle Map GQ-1248.

Van Horn, R., 1972, Surficial and bedrock geologic map of the Golden quadrangle, Jefferson County, Colorado: United States Geological Survey Miscellaneous Geologic Investigations Map I-761-A.

Wells, J.D., 1967, Geology of the Eldorado Springs quadrangle, Boulder and Jefferson Counties, Colorado: United States Geological Survey Bulletin B-1221-D.

For further information or assistance visit or call the Colorado Geological Survey at:

Colorado Geological Survey / Colorado School of Mines
1801 19th Street, Golden, CO 80401
(303) 384-2655 / CGS_Pubs@mines.edu
<http://coloradogeologicalsurvey.org/>



COLORADO GEOLOGICAL SURVEY
COLORADO SCHOOL OF MINES

