COLORADO GEOLOGICAL SURVEY
COLORADO SCHOOL OF MINES
GOLDEN, COLORADO



This mapping project was funded jointly by the Colorado Geological Survey and the U.S. Geological Survey through the National Cooperative Mapping Program under STATEMAP agreement G20AC00245.

GEOLOGIC MAP OF THE VALLEY VIEW SCHOOL QUADRANGLE WELD COUNTY, COLORADO

By Stephen M. Keller and Alexander E. Marr

2021

DESCRIPTION OF MAP UNITS

SURFICIAL DEPOSITS

For more detailed information on grain size, sorting, color, secondary calcium carbonate, and other characteristics of surficial map units see Table 1; general characteristics are presented in the text describing the map units. For details on geologic hazards (e.g., flood potential) of map units, see the Plate 2 section on Mineral Resources, Groundwater Resources, and Geologic Hazards. For optically stimulated luminescence (OSL) ages of select map units see Table 2, and for results of grain-size analyses of select units see Table 3. Subdivision of the Quaternary follows the International Chronostratigraphic Chart, v. 2020.01 (International Commission on Stratigraphy, International Union of Geological Sciences, 2020). Particle-size designation is according to the Udden-Wentworth grain-size scale (see Nichols, 2009), but (in Table 1) modified such that each sand-size category is divided into a coarser (U) and a finer (L) sub-category with the aid of a sand grain-size card (US GeoSupply, Incorporated, 2020). For example, very coarse (vc) sand is divided into vc (very coarse sand) and vcL (very coarse sand – finer).

In Table 1 the colors are designated with the aid of Munsell soil color charts (Munsell Color, 1991). Description of soil-carbonate morphology is after Machette (1985), and the type of effervescence of sediment or soil observed when treated with dilute hydrochloric acid is according to U.S. Department of Agriculture (2018).

HUMAN-MADE DEPOSITS

af — **Artificial fill (Upper Holocene)** — Mostly riprap, fill material, and, locally, refuse placed during construction of roads, railroads, buildings, dams, and landfills. Includes waste piles composed of excavated sediment adjacent to sand or gravel pits. The fill deposits generally consist of unsorted clay, silt, sand, and rock fragments. Typically, the unit is less than 6 m thick. Some deposits of artificial fill are not mapped because they are too small to show at map scale. Artificial fill may be subject to settlement, slumping, and erosion if not adequately compacted and (or) if placed on unstable slopes.

ALLUVIAL DEPOSITS

Qa — **Alluvium, undivided (Upper Holocene)** — Unit Qa covers the flat valley floor of Box Elder Creek, a north-flowing minor tributary of the South Platte River. In its physical properties and position in the landscape, unit Qa correlates with the unit Qa of Beebe Draw in the La Salle quadrangle (Palkovic and others, 2019), adjacent to the west. Beebe Draw also is a north-flowing tributary of the river. There are few exposures of unit Qa in the Valley View School quadrangle and one in the Hardin quadrangle (Palkovic and others, 2021b) adjacent to the east. One of the Valley View School quadrangle exposures is in the bank of Box Elder Creek, where the channel has been routed to run adjacent to the south side of Weld County Road (WCR) 48, 0.2 km east of WCR 57. The other two Valley View School quadrangle exposures are in the north and south ends of a sand pit located at Box Elder Creek and ~0.6 km east of the terminus of WCR 44. The Hardin quadrangle exposure is 0.4 km east of the Valley View School quadrangle's east boundary and on the east side of WCR 61, ~0.4 km north of WCR 50. Unit Qa is not sufficiently incised to form terraces along the Box Elder Creek valley. Within the active channel there are some fine-grained deposits of alluvium younger than unit Qa. These probably are correlative with alluvium one (Qa$_{1}$) of Greeley quadrangle (Keller and Morgan, 2020) and other northern Colorado Piedmont quadrangles, but are not extensive enough to be shown at map scale.

The WCR 48 exposure of unit Qa is 1.8 m thick and has the following characteristics; from ground surface downward: 0.9 m of pale-brown, as is very well sorted very fine sand with minor silt; 0.3 m of dark-brown, poorly sorted very fine to medium sand with minor silt and clay; and 0.6 m (exposed – actual thickness is greater) of grayish-brown, moderately sorted fine to medium sand with trace coarse sand. In the sand pit east of WCR 44 there are unit Qa exposures in the south and north ends of the pit. At the south end of the pit, in the east face, unit Qa overlies ~2.5 m of exposed unit Qes (on the geologic map, see sample locations V132DOSL1, V132DOSL2D2, and V132DOSL3). The unit Qa exposure is ~1.8 m high and has the following characteristics (from ground surface downward): 0.6 m of dark-brown, very cohesive clayey silt with a subordinate amount of fine to very coarse sand, no secondary calcium carbonate; 0.4 m of dark grayish-brown, very cohesive, resistant silty clay or clayey silt with a subordinate amount of fine to very coarse sand, abundant secondary calcium carbonate nodules and filaments; 0.8 m of grayish-brown, very cohesive, resistant clayey silt or silty clay, abundant secondary calcium carbonate nodules 2 to 3 cm in diameter. From the southern sand-pit exposure unit Qa can be traced southward for ~200 m to the northern sand-pit exposure, a cut bank consisting of 1.3 m of massive, very cohesive, medium dark-gray silty clay or clayey silt with abundant white calcium carbonate nodules up to 10 mm. The Hardin quadrangle exposure is a bank cut-bank along Box Elder Creek. It is 1.9 m high and is described as follows (from ground surface downward): 0.4 m of artificial fill; 0.3 m of brown, very cohesive clayey silt to very fine sand, minor medium sand; 0.3 m of dark grayish-brown, moderately cohesive sandy clayey silt, with minor very fine sand and coarse sand; 0.7 m of light grayish-brown, weakly cohesive very fine to medium sand, with minor coarse sand to granules at base; and 0.2 m (exposed) grayish-brown fine to coarse sand with fragments of carbonaceous clay.

The above exposures are the most important sources for describing unit Qa, but they are supplemented by many shallow sampling holes in this unit. These were dug with a drain spade, and some were begun at the base of road cuts and thus reached as deep as 1.3 m below undisturbed ground surface. In 40 percent of sample holes, unit Qa is silty clay or clayey silt with a fraction (5 to 35 percent) of fine to medium sand and some coarser sand. In another 40 percent, the material is very fine to medium sand with minor fines (combined silt and clay content) and minor coarse to very coarse sand. In 20 percent, the sediment is fine to very coarse sand with minor fines. In general, unit Qa is poorly to moderately sorted and moderately to very cohesive. Colors mostly are grayish-brown, dark grayish-brown, dark olive-brown, brown, and dark-brown. Secondary calcium carbonate is not prominent: in half the sample holes the alluvium has no visible calcium carbonate and no reaction to dilute hydrochloric acid; in the other half there is weak effervescence, and there are a few locations where calcium carbonate filaments or nodules are visible.

Characterization of unit Qa is assisted by lithologic descriptions in water-well driller's logs (Colorado Division of Water Resources [DWR], 2020). In the water wells used for the Quaternary cross sections (Plate 2), the alluvium is described as clay, clay with streaks of sand, fine sand and clay, and fine to medium sand. In the Box Elder Creek valley, the water-well driller's logs used for the east ends of the Quaternary cross sections indicate that unit Qa thickness ranges between 2.4 and 4.9 m, that the unit is underlain by interbedded sand, gravel, and clay layers of older alluvium considered to be alluvium three (unit Qa$_{3}$), and that the bedrock underlies unit Qa, at depths of as much as 26 m. In this minor tributary to the South Platte River, unit Qa covers unit Qa$_{3}$. Unit Qa does not occupy an incised flood plain flanked by unit Qa$_{1}$ terraces, as does unit Qa$_{1}$ (cover) of Beebe Draw in the valleys of the South Platte River and major tributaries such as the Cache la Poudre River.

In the Valley View School quadrangle, unit Qa is rarely well enough exposed for reliable stratigraphic sampling. Two of the radiocarbon ages used for unit Qa in the present report are from samples taken from the cut bank of Box Elder Creek in the Hardin quadrangle, adjacent to the east. The samples were taken from sediment near the top and near the bottom of the cut bank. The sampling location is ~1.1 km east of the Valley View School quadrangle's east boundary. Details for the samples are given in Palkovic and Morgan (2021b). The sample ages are 1,813-1,616 cal yr BP for sample V02SCI4-1, in clayey silt to very fine sand 0.4 m below ground surface, near the top of the cut bank; and 2,799-2,740 cal yr BP for sample V02SCI4-2, in fine to medium sand 1.4 m below ground surface, near the bottom of the cut bank. Within the Valley View School quadrangle and in quadrangles adjacent for OSL age determination at a sand-pit exposure, in the east pit face where unit Qa overlies unit Qes. The sampling location is shown on Plate 1. Sample V132DOSL2 was collected in unit Qa at 0.4 m below ground surface, from the highest of the three layers of unit Qa exposed there; the sample yielded an OSL age estimate of 825 +/– 45 yr. Sample V132DOSL1 was collected at 2.1 m below ground surface, from the lowest of the layers; the sample yielded an OSL age estimate of 3,640 +/– 180 yrs. Sample V132DOSL3 was collected from unit Qes immediately below the contact with unit Qa. It yielded an OSL age estimate of 3,165 +/– 290 yrs, a younger value than the age for overlying unit Qa. The younger end of the age range for unit Qes (using the error band) would be 3,455 yr. This is very close to 3,460 yr, which would be the older end of the age range for OSL sample V132DOSL2 (the younger unit Qa near the QaQes contact. Therefore, the contact may have an age of ~3,460 yr, and the OSL ages can be interpreted as indicating that the overlying unit Qa is not much younger than the underlying unit Qes. All the above ages are Late Holocene. The unit Qa OSL ages in the sand-pit face in the Valley View School quadrangle are approximately equivalent to the unit Qa radiocarbon ages in the Box Elder Creek cut bank in the Hardin quadrangle.

The correlation of map units chart (CMU) for the Valley View School quadrangle is supported by the following data sources: unit Qa and Qa$_{1}$ are not exposed in the quadrangle. Based upon relative ages of the geologic units, however, unit Qa is locally incised into unit Qe (eolian sediment) (see cross sections B-B', C-C', and D-D').

On the basis of water-well driller's logs, unit Qa is shown in cross sections as overlying unit Qa$_{1}$. Areas mapped as Qa may be prone to flooding (see Mineral Resources, Groundwater Resources, and Geologic Hazards in Plate 2).

Qa$_{1}$

Alluvium three (Upper Pleistocene) — Unit Qa$_{3}$ (also mapped as Broadway Alluvium by other researchers in the region) is mapped in the flat areas of the South Platte River valley near the northeast and northeast corners of the quadrangle. Unit at unit Qa$_{3}$ in the quadrangle, however, is inferred from subsurface water-well driller's logs (Colorado DWR, 2020). Unit Qa$_{3}$ is present in the Box Elder Creek valley, where it underlies unit Qa and overlies Laramie Formation bedrock. Unit Qa$_{3}$ in this tributary valley is not incised as it is in the South Platte River valley. Unit Qa$_{3}$ is not exposed in the quadrangle, but its subsurface extent is depicted in the cross sections (Plate 2). In the northwest corner of the quadrangle, water-well driller's logs indicate unit Qa$_{3}$ thickness as 9 m and depth to bedrock as 12 m. In cross sections B-B', C-C', and D-D' unit Qa$_{3}$ occupies a paleovalley coincident with but wider than the present Box Elder Creek valley. Unit Qa$_{3}$ thickness ranges from 11 to 26 m and depth to bedrock is between 14 and 31 m. Unit Qa$_{3}$ thickness decreases westward away from the valley axis to the (very subdued) valley wall. In cross section C-C' unit Qa$_{3}$ extends west ~0.25 km from the west wall of the present valley, and west of the present valley wall it decreases from 13 m to 4 m in thickness. In cross section B-B', west of the hill capped with unit Qg$_{2}$, a body of unit Qa$_{3}$ occupies a possible reentrant of the South Platte River valley, or a possible paleovalley that could have been a tributary to the South Platte River valley. In this area the unit Qa$_{3}$ thickness is from 3 m up to more than 8 m. Unit Qa$_{3}$ also is found in the valley of Beebe Draw (see west ends of cross sections B-B' and C-C').

In the Colorado Piedmont north of Denver, unit Qa$_{3}$ is areally and volumetrically the largest alluvium in the South Platte River valley and its major tributary valleys. This is evidenced by cross sections in Lindsey and others (2005); by Colorado Geological Survey mapping in the La Salle quadrangle adjoining the Valley View School quadrangle on the west (Palkovic and Morgan, 2019), Greeley quadrangle diagonally adjoining on the northwest (Keller and Morgan, 2020), Kersey quadrangle adjoining on the north (Lindsey and others, 2020); and geologic maps of the Masters quadrangle (Berry and others, 2015a) and Orchard quadrangle (Berry and others, 2015b), 19 to 38 km east along the South Platte River from the Valley View School quadrangle. In the northeast corner of the Valley View School quadrangle the area mapped as unit Qa$_{3}$ underlies the Kersey terrace of Bryan and Ray (1940), which flanks the south side of the South Platte River valley. From west to east the terrace extends across parts of the La Salle, Greeley, Kersey, Valley View School, Barnesville, and Hardin quadrangles. The terrace is bounded to the north by the valley floor composed of unit Qa, and by a discontinuous incised terrace of unit Qa$_{2}$. The top of the unit Qa$_{3}$ terrace is ~3 m below the top of the Qa$_{2}$ terrace (Kersey terrace).

According to Madole (2016), unit Qa$_{3}$ alluvium consists of vertical sequences of superposed, longitudinal bars of sand and gravel deposited by braided streams. These sequences locally contain finer-grained layers that probably are overbank deposits. In the water-well driller's logs (Colorado DWR, 2020) used to construct cross sections B-B', C-C', and D-D', unit Qa$_{3}$ consists of interbedded clay, sand, and gravelly sand-sandy gravel layers in approximately equal proportions. Gravel layers typically are present toward the bottom of the unit and clayey layers toward the top. In the Greeley quadrangle (Keller and Morgan, 2020) layers of unit Qa$_{3}$ in the valley of the South Platte River and Cache la Poudre River are typically composed of: 1) gravelly sand and sandy gravel (preponderant); 2) very fine to very coarse sand with little or no gravel (subordinate); and 3) clayey silt to clay with minor gravel, possibly overbank deposits (rare). Gravel-pit faces in the Greeley quadrangle expose the following: interbedded sand and gravel layers, as well as gravel-filled channels inset in underlying sand and gravel layers; crossbedding in gravel layers; and large (as much as 0.2 m thick and 1.5 m long) rip-up fragments of sand layers contained in gravel layers (Keller and Morgan, 2020). Also in the Greeley quadrangle, three south-trending tributary paleovalleys join the Cache la Poudre River valley on the north side, and in these valleys (as seen on hillshaded lidar imagery) and subsurface dimensions they resemble the paleovalley of Box Elder Creek in the Valley View School quadrangle. Based on a Greeley quadrangle cross section through Quaternary units, unit Qa$_{3}$ sediments in the tributary Sand Creek paleovalley are continuous and coeval with unit Qa$_{3}$ sediments in the Cache la Poudre River valley (Keller and Morgan, 2020). By analogy it is probable that unit Qa$_{3}$ in the Box Elder Creek paleovalley is continuous and coeval with unit Qa$_{3}$ in the South Platte River valley.

There are no unit Qa$_{3}$ exposures in the quadrangle and thus no samples could be obtained for laboratory age determination. Deposits of unit Qa$_{3}$ have been dated by the CGS in several nearby quadrangles, such as the Greeley and Gowanda quadrangles (Keller and Morgan, 2020; Keller and others, 2019), and have been assigned to the Late Pleistocene. In the Valley View School quadrangle, unit Qa$_{3}$ locally is overlain by Early to Middle Holocene eolian sediment (unit Qe). Late Holocene eolian sand (unit Qes), and Late Holocene alluvium (undivided unit Qa) (see cross sections B-B', C-C', and D-D'). Coal seams recorded in water-well driller's logs indicate that the sandstone bedrock beneath unit Qa$_{3}$ is the lower part of the Laramie Formation. In the Box Elder Creek valley unit Qa$_{3}$ is an important shallow alluvial aquifer and its sand and gravel layers are possible sources of aggregate resources.

Qa$_{3}$

Gravel one (Middle Pleistocene) — Unit Qg$_{1}$ caps two hills in the northeast corner of the quadrangle. The hills are areas of elevated topography in that the capping gravel is more resistant than the underlying bedrock. Unit Qg$_{1}$ characteristics are described mostly from a shallow trench (part of this mapping project) excavated at ground surface near the center of the northernmost hill. The trench includes OSL sampling locations V017COSL1 and V017COSL2, as shown on the Plate 1 geologic map. Other unit Qg$_{1}$ data from the hill are from an inactive former gravel pit adjacent to the north of and several meters lower in elevation than the trench, from gravel inferred at a probe dug low-on-the-face of the trench, and from a road cut on WCR 57. The mapped areal extent of the unit is inferred based upon landforms visible in the lidar image for the quadrangle, the deposit thickness as estimated using local topography, and the National Resource Conservation Service (NRCS) soils map of the quadrangle (Crabb, 1980). At the hill, the thickness of unit Qg$_{1}$ is ~5 m. Approximately 1 m of the base of the unit is exposed in the adjacent former gravel pit and overlies Laramie Formation sandstone. The trench was dug entirely in unit Qg$_{1}$, and the exposed trench face was 2.2 m high; the bottom of the face was ~3 m above the Laramie Formation.

In the sampling trench, unit Qg$_{1}$ consisted of the following material (from ground surface downward): 0.4 m of weakly cohesive, poorly sorted gravelly sand, clasts up to 6 cm, brown, some secondary calcium carbonate; 0.5 m of moderately cohesive sandy gravel, clasts up to 7 cm with calcium carbonate rinds coating the bottoms of many clasts, white, matrix locally cemented by calcium carbonate; 0.6 m of weakly cohesive, poorly sorted medium to very coarse sand with minor gravel, some secondary calcium carbonate; 0.6 m of strongly cohesive, poorly sorted gravelly sand or sandy gravel, clasts up to 11 cm with calcium carbonate rinds, pinkish-gray, matrix locally cemented by calcium carbonate; and 0.1 m of weakly cohesive, well sorted medium to coarse sand with no fines or gravel, pinkish-gray, no secondary calcium carbonate. In the gravel pit exposure, unit Qg$_{1}$ is a poorly sorted, strongly cohesive gravelly sand with clasts up to 17 cm, pale-yellow, and with abundant secondary calcium carbonate (Stage III) both as matrix and nodules.

In the Valley View School quadrangle, the highest surface of unit Qg$_{1}$ is ~40 m above the South Platte River, which is ~4 km to the northeast. In the Masters quadrangle (Berry and others, 2015a), ~28 km southeast of the Valley View School quadrangle, three exposures of unit Qa (intermediate alluvium) are in a long line ~2 km long and parallel to the South Platte River; the river lies 2.5 km to the north, and the tops of the exposures are 30 to 36 m above the river. Unit Qai of the Masters quadrangle and unit Qg$_{2}$ in the Valley View School quadrangle both have pervasive secondary calcium carbonate in the matrix and carbonate rinds on clasts. The gravel remnants in the above two quadrangles are at a similar distance southwest of the South Platte River. Based on uranium-series dating, Berry and others (2015a) determined the best unit Qai numerical age estimate as somewhat older than 382 ka. They correlate the unit with older deposits of Slocum Alluvium, per the age range of 320-390 ky for this unit as given in Kellogg and others (2008).

Unit Qg$_{1}$ trench sample V017COSL1, from sandy gravel 1.1 m below ground surface, yielded an age of >80,000 yr from conventional OSL analysis; unit Qg$_{1}$ trench sample V017COSL2, from sandy gravel 2.0 m below ground surface, yielded an age of >90,000 from conventional OSL analysis (Table 2). Considering the advanced secondary carbonate accumulation in the gravel, and relatively high elevation of the deposit, these ages were unexpectedly young. A second analysis of unit Qg$_{1}$ trench sample V017COSL2 yielded an age of 341,080+/-22,840 yr from thermal transfer OSL (TT-OSL) analysis (Table 2). This age is not greatly younger than the 382 ka age for unit Qai and also falls within the Slocum Alluvium age range. Therefore, unit Qg$_{1}$ in the Valley View School probably is equivalent to unit Qai in the Masters quadrangle and also correlative with Slocum Alluvium as mapped elsewhere in the Colorado Piedmont.

Qg$_{1}$

EOLIAN DEPOSITS

Qes — **Eolian sand (Upper Holocene)** — Unit Qes covers approximately half the total area and much of the upland in the Valley View School quadrangle. It lies within an eolian sand area named the "South Platte deposit" by Madole and others (2005). This sand body is one of several extensive, eastern Colorado eolian sand deposits mapped and described in that publication. In the field, unit Qes is recognized in road cuts and in sample holes that penetrate below in thin (generally less than 20 cm thick) and poorly developed soil A horizon (for definitions of soil horizons see Birkeland, 1999). Unit Qes eolian sand is distinguishable from unit Qe eolian sediment because the former consists of coarser, well-sorted sand with only a small amount (less than 6%) of fines (silt and clay), whereas the latter consists of finer, less-sorted sand locally containing a large amount (22 to 30%) of fines. More evidence supporting this distinction is presented by Keller and Morgan (2020). Mapping of unit Qes in the Valley View School quadrangle is supported by the following data sources: road cuts, excavations, many shallow sample holes, 121 water-well driller's logs (Colorado DWR, 2020), the extent of eolian sand as mapped by Madole and others (2005), the distinctive northwest-to-southeast-oriented dune topography of unit Qes as recognized in the lidar image for the quadrangle, and the NRCS soils map of the quadrangle (Crabb, 1980). Within the South Platte sand deposit (Madole and others, 2005), on the west side of the Valley View School quadrangle, there is a group of seven "windows" or gaps in the eolian sand cover; these areas are roughly 1 to 3 km in diameter. Water-well driller's logs, sample holes, and the NRCS soils map for the quadrangle indicate that these areas mostly consist of Laramie Formation (see geologic map). Also, west of the Box Elder Creek valley there are some areas included in the eolian sand cover by Madole and others (2005) but which are designated as unit Qe on the present map. This distinction is based on water-well driller's logs, sample holes, and the soils map. (It should be noted that Madole and others (2005) is a small-scale reconnaissance map covering a very large area.) In the northern third and middle third of the quadrangle, represented by cross sections B-B' and C-C' respectively, water-well driller's logs in the cross sections indicate unit Qes thickness to be in the range of 4 to 8 m. In the southern third, cross section D-D' depicts the thickness as mostly 4 to 10 m. All three cross sections show unit Qes to be thicker closer to hilltops and thinner on hillslopes and in valleys. On the basis of lithology and position in the landscape, unit Qes correlates with eolian sand mapped by the CGS in other Colorado Piedmont 7.5' quadrangles, such as Frederick (Keller and Morgan, 2018), La Salle (Palkovic and others, 2019), and Greeley (Keller and Morgan, 2020). In the Valley View School quadrangle there are no exposures of unit Qes in contact with underlying deposits. On the basis of water-well driller's logs (Colorado DWR, 2020), unit Qes locally overlies Laramie Formation bedrock, alluvium of unit Qa$_{3}$, and eolian sediment of unit Qe (see cross sections B-B', C-C', and D-D'). In the Frederick quadrangle, 42 km to the southwest of the map area, unit Qes was observed overlying unit Qe in a shallow trench and also in nearby geotechnical boreholes (Keller and Morgan, 2018).

Unit Qes is composed of poorly sorted very fine to coarse or very coarse sand, moderately sorted fine to medium sand with minor coarse to very coarse sand, and poorly sorted fine to coarse sand with minor very coarse sand. Fines are minimal at all locations and rare granules and small pebbles are present at one-third of the sample locations. At sample V132DOSL3 (see geologic map) one sample was taken from unit Qes for grain size analysis. The material is composed of 94.4% sand, 2.6% silt, and 3.0% clay (Table 3), which is in accordance with the low content of fines (4 to 9%) reported for eolian sand by Madole and others (2005). The sand is non-cohesive to weakly cohesive, lacks visible secondary carbonate, and has no reaction to dilute hydrochloric acid. Colors are brown, yellowish brown, grayish brown, and pale brown. Unit Qes is best observed in the east and west faces of a sand pit located at Box Elder Creek and ~0.6 km east of the terminus of WCR 44 (location of grain size sample V132DSGS, above), and in a smaller sand pit located ~0.5 km to the north. The deposits exposed in these pits are planar bedded at a scale of several millimeters to several centimeters, locally cross-bedded, and have stoss-side dune beds with a northward apparent dip of ~20 degrees. At the sand pit, OSL sample V132DOSL3 was collected at 1.9 m below ground surface, at the top of unit Qes and within a few cm below the contact with overlying unit Qa. This sample yielded an age of 3,165+/-290 yr, which is Late Holocene and accords with the Madole and others (2005) Late Holocene age group for Qes (the youngest of three Qes age groups presented in that publication). Unit Qes sand has been extracted (possibly for construction material) from small excavations east of the Box Elder Creek Valley. There is abundant, relatively clean eolian sand remaining at these sites and elsewhere in the quadrangle. The area mapped as unit Qes locally may contain collapsible soils.

Qe — **Eolian sediment (Lower Holocene to Upper Pleistocene)** — Eolian sediment of unit Qe covers approximately one-third of the upland area west of the Box Elder Creek valley. This deposit occupies much of the north part of the Valley View School quadrangle and extends northward into the Kersey quadrangle. Less-extensive deposits are present in the center and south of the quadrangle. Mapping of unit Qe in the Valley View School quadrangle is supported by observations in road cuts, sample holes, grain-size analysis, water-well driller's logs (Colorado DWR, 2020), and the NRCS soils map for the quadrangle (Crabb, 1980). At present, unit Qe occupies much less of the quadrangle's ground surface than does the eolian sand of unit Qes. Cross sections B-B', C-C', and D-D' indicate, however, that Qe is not associated with particular landforms and can be present over a broad range of elevations. On the basis of its position in the landscape, lithologic characteristics, and secondary calcium carbonate development, unit Qe is similar to unit Qe in the Longmont quadrangle (Madole, 2016), Milliken (Palkovic and others, 2018), Greeley quadrangle (Keller and Morgan, 2020), and other Colorado Piedmont quadrangles mapped by the CGS. In the northern third and middle third of the Valley View School quadrangle, as represented by cross sections B-B' and C-C' respectively, water-well driller's logs indicate that unit Qe thickness is typically 2 to 7 m. In the southwest corner of the quadrangle, the west end of cross section D-D', unit Qe has a thickness range of 9 to 14 m. Unit Qe locally may contain small unmapped bodies of unit Qes.

Unit Qe is composed of poorly sorted, very fine sand to coarse sand, locally with a dominant fine fraction, and poorly sorted fine to medium sand with minor coarse to very coarse sand, also locally with a dominant fines fraction. At a few locations the deposit contains rare to more granules and small pebbles. The fines fraction is usually sufficient to impart moderate cohesion, although cohesion locally can be either weak or strong. Unit Qe is massive in the two road cuts where it is exposed (the deposit also is massive in excavations in other piedmont quadrangles mapped by the CGS). Colors are brown, grayish-brown, dark grayish-brown, and dark yellowish-brown. Unlike in other quadrangles mapped by the CGS, unit Qe sampled in the Valley View School quadrangle has no visible secondary calcium carbonate and no reaction to dilute hydrochloric acid. This may be because road cuts and sample holes did not penetrate deep enough to encounter secondary calcium carbonate (e.g., in a soil Bk horizon). Two sediment samples were taken from unit Qe for grain-size analysis; the locations are V030GS and V035OS on the geologic map. Their respective sand fractions are 77.7 and 68.8%, silt fractions 14.4 and 17.5%, and clay fractions 7.5 and 12.7% (Table 3). The total fines in the two samples were 21.9 and 30.2%, respectively, which is in accord with total fines being mostly between 21 and 55% in 19 geotechnical grain size analyses of unit Qe in the Greeley quadrangle, adjacent diagonally to the northwest (Keller and Morgan, 2020). The unit Qe silt fraction in the Valley View School and Greeley quadrangles is too small for the material to be considered sandy loess.

On the basis of water-well driller's logs (Colorado DWR, 2020), unit Qe overlies Laramie Formation bedrock and the alluvium of unit Qa$_{3}$; it locally underlies the eolian unit of unit Qes (see cross sections B-B', C-C', and D-D'). Along the west side of the Box Elder Creek valley, unit Qe is interpreted to be incised by a stream that deposited alluvium of unit Qa. In the Frederick quadrangle, 42 km to the southwest, unit Qe was observed underlying unit Qes in a shallow trench and also in nearby geotechnical boreholes (Keller and Morgan, 2018).

In the Valley View School quadrangle, unit Qe is not sufficiently exposed to collect suitable samples for radiocarbon analysis. Radiocarbon ages for unit Qe are ~9 ky and ~21 ky in the Johnstown quadrangle (Palkovic and Morgan, 2017), ~4 ky and ~5 ky in Frederick (Keller and Morgan, 2018), ~5 ky in Gowanda (Keller and others, 2021), and ~9 ky in Greeley (Keller and Morgan, 2020). With one exception (~21 ky age in the Johnstown quad) these ages are Holocene. Within the Valley View School quadrangle, unit Qe is not a likely potential source of clean sand because of its relatively high content of fines. The area mapped as unit Qe locally may contain collapsible soils.

BEDROCK GEOLOGY

Within the Valley View School quadrangle, the only bedrock unit mapped at the ground surface is the Laramie Formation, and its mapping is mostly limited to residium formed in place by weathering of the formation. Bedrock units beneath the Laramie Formation are shown in cross section A-A' (Plate 2). Descriptions of the Laramie Formation and Fox Hills Formation are from Spencer (1986), who investigated these units in detail in the Frederick and Erie quadrangles, ~46 km southwest of the Valley View School quadrangle. Descriptions of Pierre Shale members are adapted from Scott and Cobban (1965), supplemented by descriptions by Braddock and others (1988). Descriptions of older units are adapted from descriptions in the geologic map of the Carter Lake Reservoir quadrangle, 53 km to the west, where these units are described in outcrop (Braddock and others, 1988). Thickness values for the Laramie Formation and Fox Hills Sandstone are from Spencer (1986). Thickness values for older bedrock units are from cross section A-A', which was prepared with the aid of oil and gas well logs from the Colorado Oil and Gas Conservation Commission (COGCC) (2020).

Ki — **Laramie Formation (Upper Cretaceous)** — The Laramie Formation is mostly overlain by resiudum developed mainly from the weathering of sandstone, and most of this residuum is on the upland in the west side of the quadrangle. There is one fresh exposure of Laramie Formation sandstone, in the northeast corner of the quadrangle and in an area not accessible on a gravel pit adjacent to the north of OSL sample locations V017COSL1 and V017COSL2. Also, fragments of sandstone with small (<6 cm in diameter) zones of carbonaceous mineral were found beneath unit Qa and unit Qes, at the base of a shallow trench located near the quadrangle's west boundary and near where the Gilmore Ditch crosses WCR 48. Laramie Formation residuum is mapped from sample holes, water-well driller's logs (Colorado DWR, 2020), the NRCS soils map of the quadrangle (Crabb, 1980), and areas of non-deposition of the South Platte eolian sand area mapped by Madole and others (2005). At the gravel pit mentioned above, the Laramie Formation exposure is ~1 m high and consists of well-sorted, very fine-grained to fine-grained sandstone with trace medium sand and minor fines. The sandstone is weakly to moderately cohesive and pale-yellow, and has planar and flexed bedding at a scale of millimeters to 2 cm. It is overlain by 0.5 m of unit Qg$_{1}$ gravelly sand. In the Erie and Frederick quadrangles, ~50 km southwest of the map area, the lower 60 to 90 m of the Laramie Formation consists of light- to medium-gray quartzose sandstone layers separated by clay, fire clay (refractory clay), shale, or coal seams. The upper 123 to 153 m consists of claystone, shale, sandy shale, and lenticular beds of sandstone and lignite. Total thickness is 213 m. Multiple coal beds are present in the lower part of the formation and are as much as 3.7 m thick (Spencer, 1986). Coal seams reported in water-well driller's logs in the Valley View School quadrangle indicate that the lower part of the Laramie Formation underlies the Quaternary deposits in the quadrangle. At the south end of cross section A-A' (Plate 2), limited data for the Laramie Formation indicate that the coal and gas wells penetrate as much as 76 m of the unit (COGCC, 2020). Data used in Quaternary cross sections B-B', C-C', and D-D' (Plate 2) indicate that water wells penetrate up to 79 m of the formation (Colorado DWR, 2020).

Kfh — **Fox Hills Sandstone (Upper Cretaceous)** — Greenish-tan, fine- to coarse-grained quartzose sandstone; cross bedded in the lower part, grading upward to a light-yellow and white, massive, fine- to medium-grained sandstone. Thickness is 46 to 91 m. The contact of the Fox Hills Sandstone with the underlying Pierre Shale is not depicted on cross section A-A' because data for this contact are not included in the oil and gas well logs used for the cross section.

Pierre Shale (Upper Cretaceous) — Marine strata composed of dark-gray shale, siltstone, and fine-grained sandstone. Bentonite beds, rich in weathered volcanic ash, are common in its lower part and calcareous concretions are common throughout. The various members of the Pierre Shale contain index fossil ammonite species (Scott and Cobban, 1965).

Kpk/Kpu — **Upper most transition member and upper shale member, undivided** — These two members are combined in cross section A-A' because they are not distinguished in the oil and gas well data used to prepare the cross section. Both members are observed in outcrop ~45 km southwest of the Valley View School quadrangle (Scott and Cobban, 1965). In general, the members consist of friable sandstone and soft shaly sandstone containing thin-bedded sandy shale and calcareous sandstone concretions. Combined thickness of the two members is ~1,060 m.

Kpl — **Larimer and Rocky Ridge Sandstone Members and intervening shale, undivided** — The Larimer and Rocky Ridge Sandstone Members are well-indurated, light-gray to light-brown, medium-grained sandstone, composed mostly of quartz with minor feldspar and biotite. Combined thickness is ~46 m.

Kpm — **Middle shale member** — This unit consists of highly friable, greenish-gray claystone and sandy siltstone, and contains bentonite beds. Thickness is ~120 to ~220 m.

Kpn — **Hygiene Sandstone member, undivided** — The upper part of the unit is well-indurated, light-gray, fine- to medium-grained sandstone composed mostly of quartz, minor feldspar, and minor opaque minerals. The middle part of the unit is medium-gray siltstone; the lower part is a friable, gray concretionary sandstone. Thickness is ~185 to ~200 m.

Kpl — **Lower shale member** — Dark olive-gray shale and sandy shale containing limestone and ironstone concretions; bentonite beds are common in the lower part of the unit. Thickness is ~470 to ~540 m.

Kn — **Niobrara Formation, undivided (Upper Cretaceous)** — Very fissile, dark-gray shale containing thin (5 m) micritic limestone layers. Unit is an important oil and gas source in the quadrangle and in much of the Denver Basin (see Plate 2 and section on Mineral Resources, Geosystems, v. 16, p. 2508–2528 [also available at https://doi.org/10.1002/2015GC005881].

Kcgg — **Colorado Group - Carlile Shale, Greenhorn Limestone, Graneros Shale, and Mowry Shale (undivided) (Upper Cretaceous)** — Olive-gray silty claystone and sandy claystone; dark-gray limestone and olive-gray calcareous, silty claystone and siltstone; dark-gray to grayish-black siltstone and claystone and siliceous shale. Combined thickness is ~140 m.

Kd — **Dakota Group: South Platte Formation and Lytle Formation (undivided) (Lower Cretaceous)** — Unit consists of gray to light-brown, well-sorted, fine- to medium-grained sandstone; dark-gray carbonaceous shale; gray to light-brown, fine-grained sandstone; and gray to light-brown, coarse-grained, conglomeratic sandstone. Combined thickness is ~155 m. A maximum of 36 m of this unit was penetrated in the two oil and gas wells used to prepare cross section A-A'.

MAP SYMBOLS

Contact — Approximately located

Normal fault of Weimer (1996) — Existence certain, location concealed, where known, sense of displacement indicated by ball and bar on downthrown side

Longmont wrench fault zone — Existence certain, location concealed; ball and bar on downthrown side

Strike-slip fault (in cross section) — A, away from observer; T, toward observer

Oil and gas wells

Water well

Soil boring for geotechnical investigation

Grain-size analysis sample site

Optically stimulated luminescence (OSL) sample site

Dune crest

Eolian blowout

Gravel pit — Sand or gravel pit - exposed sand or gravel faces

A – A' Alignment of cross section

REFERENCES